

Medication Information

Effexor

Deplin is a "medical food" used in people with depressive to enhance the effects of antidepressant medications.

You should not use this medication if you have ever had an allergic reaction to I-methylfolate. If you have certain conditions, you may need a dose adjustment or special tests to safely use this medication. Before using I-methylfolate, tell your doctor if you have:

- a history of bipolar disorder (manic depression);
- vitamin B12 deficiency; or
- pernicious anemia.
- are pregnant

Avoid drinking alcohol while you are taking I-methylfolate.

Get emergency medical help if you have any of these signs of an allergic reaction: hives; difficulty breathing; swelling of your face, lips, tongue, or throat.

Less serious side effects may include:

- nausea
- loss of appetite
- bloating gas
- bitter taste in your mouth.

Drugs that can interact with I-methylfolate include:

- birth control pills
- cholestyramine (Prevalite, Questran), colestipol (Colestid)
- colchicine
- isotretinoin (Accutane)

methylprednisolone (Medrol)

- methotrexate (Rheumatrex, Trexall)

pancreatin (Hi-Vegi-Lip), pancrelipase (Cotazym, Pancrease, Ultrase)

- pentamidine (Nebupent)

pyrimethamine (Daraprim)

sulfasalazine (Azulfidine)

- triamterene (Dyrenium)

- trimethoprim (Proloprim, Bactrim, Septra) •

non-steroidal anti-inflammatory medication (NSAIDs) most seizure medications